

RESEARCH ARTICLE

Hareport hazard: Identifying hare activity patterns and increased mammal–aircraft strike risk at an International Airport

Samantha Ball^{1,2} , Anthony Caravaggi³  & Fidelma Butler¹ ¹School of Biological, Earth and Environmental Science, Distillery Fields, University College Cork, Cork T23 TK30, Ireland²daa, Airport Fire & Rescue Service, Dublin K67 CX65, Ireland³School of Applied Sciences, University of South Wales, 9 Graig Fach, Glyntaff, Pontypridd CF37 4BB, UK**Keywords**

Airfield management, circadian activity, wildlife hazard, wildlife management, wildlife strikes

Correspondence

Samantha Ball, School of Biological, Earth and Environmental Science, Distillery Fields, University College Cork, Cork T23 TK30, Ireland. Tel: +353 21 4904676; Email: samantha.ball@ucc.ie

Editor: Marcus Rowcliffe

Associate Editor: Oliver Wearn

Funding Information

daa: EBPPG/2018/43 Irish Research Council University College Cork Consortia

Received: 1 March 2022; Revised: 10 June 2022; Accepted: 27 June 2022

doi: 10.1002/rse2.293

Abstract

Reported strike events between wildlife and aircraft are hazardous to aircraft and airfield operations and are increasing globally. To develop effective mitigation strategies, the relative hazard a species poses to aircraft, as well as information relating to its life history, are key to the development of effective mitigation strategies in Wildlife Hazard Management Plans. However, given the complex nature of airfield environments with access restrictions and the presence of sensitive equipment, the collection of high-quality ecological data can be difficult. Here we use motion-activated camera traps to collect activity data on a population of Irish hares (*Lepus timidus hibernicus*) inhabiting the airfield at Dublin International Airport, to investigate the link between hare activity and aircraft activity in relation to hare strikes. Camera traps revealed that the hare population at the airfield largely displayed a bimodal crepuscular activity pattern, with activity peaking at sunrise and at sunset. Recorded hare strike times at the airfield were closely associated with hare activity times with a high temporal overlap between these datasets. In comparison, hare activity and aircraft movement activity had a moderate overlap across all seasons, with strikes peaking at times with low aircraft movements. We demonstrate the importance of understanding the circadian and seasonal activity patterns of hazardous species at airfields for targeted strike mitigation.

Introduction

Wildlife collisions or ‘strikes’ with transport vehicles can have serious consequences for passenger safety, industry economics (e.g. Dolbeer & Begier, 2021), the local economy (e.g. Jaren et al., 1991) and wildlife conservation efforts (e.g. Clair et al., 2019). While the majority of strike-related research has focused on road traffic networks (e.g. Popp & Boyle, 2017; Wright et al., 2020), similar consequences are also reported for other modes of transport including rail (Dorsey et al., 2017), shipping networks (Laist et al., 2001) and air transportation (Altringer et al., 2021). Mammals are well represented within the literature regarding strikes for most modes of transport (e.g. Pokorný et al., 2022), yet relatively little research has focussed on mammals in the context of the

air transportation sector, despite mammalian strikes composing 3–10% of reported strikes in the aviation industry (Ball, Caravaggi, & Butler, 2021b). Terrestrial mammals are hazardous to aircraft only when they move on to the active runway, therefore, understanding the circadian (over 24 hours) and seasonal activities of animals inhabiting or using the airfield could help to identify periods of increased risk when animals are likely to come into contact with aircraft. Identifying these periods of risk can then allow for the targeted development and application of strike mitigation measures.

Animal behaviours and activity patterns are greatly influenced by a variety of pressures within their environment, including- but not limited to- food availability (Pereira, 2010), predation risk (Ross et al., 2013), disturbance and anthropogenic activities (Lendrum

et al., 2017). Circadian and seasonal activity patterns are photoresponsive cycles regulated by the suprachiasmatic nucleus of the hypothalamus in mammals (Meijer et al., 2010), allowing for species to fulfil daily requirements (e.g. feeding) while adapting to seasonal day length. This plasticity (Phillips et al., 2013) allows for mammals to exploit the ecological environment they inhabit and alter activities relative to changing sunrise and sunset times. Hence, in the context of wildlife management, wildlife hazards are likely to exhibit temporal variability, particularly with increasing distances from the equator where seasonality becomes increasingly pronounced. This highlights the importance of understanding species composition and associated life histories of potentially hazardous fauna on an airfield-specific basis.

Given the sensitive nature of airfield environments, the collection of high-quality ecological data can often be complicated, requiring the use of remote field methods due to limited accessibility, such as camera traps (Carswell et al., 2021; Scheideman et al., 2017), radio telemetry (York et al., 2000), GPS tracking (Askren et al., 2019) and predictive modelling based on pre-existing movement data (Arrondo et al., 2021). Lagomorphs (particularly rabbits and hares) are frequently reported in airfield environments and are reportedly involved in strike events near globally (Ball, Caravaggi, & Butler, 2021b; Dolbeer & Begier, 2021; Kitowski, 2016). A population of the Irish hare (*Lepus timidus hibernicus*, Bell 1837), an endemic subspecies of the Mountain hare (*L. timidus*, Linnaeus 1758), resides at Dublin Airport in the Republic of Ireland where strike events between hares and aircraft have been increasing by an average of 14% annually since 1997 (Ball, Butler, et al., 2021a). The damage potential of a hare strike (10,576 J; Ball, Butler, et al., 2021a), in tandem with the conservation status (Caravaggi et al., 2017; Reid et al., 2010) of this endemic subspecies, require that effective management strategies be developed to mitigate against strike events. Here, we investigate whether motion-activated camera traps – an easily accessible and relatively inexpensive method of monitoring – can be successfully used to identify periods of increased strike risk between aircraft and hares.

Understanding the relationship between animal activity patterns and temporal distributions of air traffic can help to comprehend and mitigate strike risk (Arrondo et al., 2021; Carswell et al., 2021; Schwarz et al., 2014). Here, we apply an approach more frequently used in inter-specific competition and predator–prey modelling (e.g. Caravaggi et al., 2018; Ross et al., 2013) to an industry setting. We hypothesize that aircraft-hare activity overlap will fluctuate seasonally, and that hare activity represents a better indicator of strike patterns than aircraft activity. We focus on the Irish hare as a model species and propose that this approach could be used to

identify periods of risk associated with other ground dwelling species at airfields worldwide and for use on public road networks.

Methods

Study area

Dublin International Airport (53.4264°N, 6.2499°W) is Ireland's largest civil airport and one of the busiest in Europe, with almost 250,000 aircraft movements recorded in 2019 alone. The airfield was composed of approximately 275 hectares (680 acres) of grassland throughout the study period, which increased to approximately 370 hectares (914 acres) in August 2021 due to the expansion of the airfield to incorporate an additional runway. The grasslands are maintained according to a long grass management policy (UKCAA Safety Regulation Group CAP, 2008) comprising of a blend of Italian ryegrass (*Lolium multiflorum*) and tall fescue (*Festuca arundinacea*). The airfield is located on the east coast of Ireland and experiences a temperate climate. A mean temperature of 9.4°C and mean rainfall of 62.8 mm was recorded throughout the study period (July 2019–May 2021; MET Eireann, 2021).

Strike data

A database of all strike events at the airfield has been maintained since 1990, encompassing all confirmed strike events with avian and mammal species. The first strike event with a hare was reported at the airfield in 1997. These data were provided by the daa (Dublin Airport's managing body). Carcasses were recovered from active areas (i.e. runways, taxiways) following a reported strike event or during mandatory routine inspections with the location and environmental conditions surrounding an event recorded (e.g. weather; see Ball, Butler, et al., 2021a). From 2012 onwards, temporal data detailing the date and time of a strike were also recorded, resulting in $n = 238$ hare strike events with an associated strike time from 2012 until December 2021. Despite the presence of other mammal species occasionally reported at the airfield including foxes (*Vulpes vulpes*), hedgehogs (*Erinaceus europaeus*), rabbits (*Oryctolagus cuniculus*), rats (*Rattus norvegicus*), domestic cats (*Felis catus*) and bats (Kelly et al., 2017), strike events with these species are rare (Bolger & Kelly, 2008). With an increasing number of hare strike events reported annually and a sufficient kinetic energy to cause damage to an aircraft (Ball, Butler, et al., 2021a), the Irish hare is the most hazardous mammal species at the airfield. Therefore, here we focus on the Irish hare due to a high incidence of strike events.

Camera trapping

Seven Bushnell Trophy Cam HD (model 119476) camera traps were deployed into the grasslands surrounding the main runway at Dublin Airport (RW28-10 L), at seven fixed locations, identified *a priori* by airfield safety authorities and dependant on the location of appropriate permanent structures (e.g., gate posts). Camera positions were not changed due to ongoing construction work throughout the study period. Cameras were positioned 50 cm above ground level with a $\sim 15^\circ$ downwards tilt with cameras pointing away from areas of high aircraft and vehicle traffic, to prevent false triggers and operated 24/7 with the use of infrared for nocturnal image capture (Appendix 1 and Appendix 2). The shortest distance between the two closest camera traps was 520 m (0.5 km, range 520 m–920 m), to minimize spatial replication. This was well beyond the typical hare home range ($0.14\text{km}^2 \pm 0.02$ m Caravaggi et al., 2016) and slightly over the median home range size of male hares reported by Wolfe and Hayden (1996) of 0.5 km^2 . Cameras were left *in situ* for 32–69 days (dependant on logistics, camera performance and accessibility), for each season between July 2019–May 2021 (Appendix 3), for a total of 360 calendar days. Seasons were defined as spring (March–May), summer (June–August), autumn (September–November) and winter (December–February). Interannual survey data were pooled for each season. Cameras were programmed to record two-time stamped images when triggered by movement at medium sensitivity, with a 60-second interval between images.

Circadian activity- data analysis

Camera trap images were assumed independent of each other if they were separated by a minimum of 30 minutes, or by a clearly different animal (e.g. distinguishable markings; Viviano et al., 2021) and therefore defined as an independent mammal detection. As camera traps were only triggered by movement (i.e., activity), we also assumed that detections of hares were a true reflection of the circadian activity of the species. For trigger events where the image contained more than one individual ($n = 39$ events), only a single event was recorded (Caravaggi et al., 2018). Aircraft movement data for the runway (RW28-10 L) were obtained from the daa, whereby a movement is defined as a take-off or landing manoeuvre. All aircraft manoeuvres were considered to be independent events. These data were truncated to include only those dates where camera traps were deployed on the airfield. All statistical analysis was carried out in programme R v 4.0.4. (R Core Team, 2021).

A cross-correlation function (CCF) was used to determine negative ($h-$) or positive ($h+$) lags in time series data between aircraft and hares. CCFs show correlations between events in time series data (X_a , Y_a ; $a = \text{time}$), where a positive lag ($h+$) shows a correlation between X_{a+i} and Y_a (i.e., X succeeds Y). Likewise, a negative lag ($h-$) shows a correlation between X_{a-i} and Y_a (i.e., X precedes Y). The significance of the correlation coefficient was established by calculating the t value, where the critical t value ($P = 0.05$, 22 degrees of freedom, one-tailed) = 1.72. The 'Overlap' package (Meredith & Ridout, 2021) was used to determine the temporal overlap between hares and aircraft, by estimating the overlap coefficient (Δ), where $\Delta = 0$ indicated no overlap and $\Delta = 1$ indicated complete temporal overlap. Data were bootstrapped 1,000 times to generate 95% confidence intervals (CI) of the overlap coefficient (Zanni et al., 2021). The Δ_4 estimator was used for all pairwise comparisons between aircraft movements and hare activity and was also used to compare aircraft movements and hare strike events across the whole year. The Δ_1 estimator was used for seasonal aircraft strike pairwise combinations due to seasonal strike records ranging from 54–69 events (Meredith & Ridout, 2021) and for seasonal hare activity vs. seasonal hare strike comparisons. Temporal overlaps between hares and aircraft were ranked as either high ($\Delta > 0.75$), moderate ($0.50 < \Delta < 0.75$) or low ($\Delta < 0.50$), for each season (Monterroso et al., 2014).

Given Ireland's northern latitude, day length varies substantially with seasons with the longest days in the summer experiencing ~ 17 hours of daylight, and the shortest days in the winter experiencing ~ 7 hours of daylight. Therefore, to investigate how activity patterns changed in relation to the rising of a setting sun seasonally, trigger events were offset to either sunrise or sunset (i.e. 00:00–11:59 offset relative to sunrise, 12:00–23:59 offset relative to sunset; Caravaggi et al., 2018). All daytime offsets (i.e., events which occurred between sunrise and sunset) were converted to positive integers and night-time offsets (i.e., events which occurred between sunset and sunrise) to negative integers. As an example, a detection at 10:15 on a day where sunrise was at 08:00 would have an offset value of +2 hrs 15mins, which would indicate diurnal activity. Similarly, a detection at 20:30 on a day where sunset was at 19:00 would have an offset value of -1 hr. 30 mins, indicating nocturnal activity. Finally, a detection which occurred at 03:55 (after midnight) would be offset to sunrise and on a day where this was at 07:15, an offset value of -3 hrs 20mins would be allocated, indicating nocturnal activity closer to sunrise than sunset. These offset values were used as the dependant variable in a one-way analysis of variance with *post hoc* Tukey tests, to test differences in activity patterns across seasons. Likewise, as

the Covid-19 pandemic substantially impacted on the number of aircraft movements at Dublin Airport, the same method was used to compare hare activity between seasons (*i.e.*, Summer 2019 vs. Summer 2020) to test whether hares exhibited altered activity patterns with the relief in aircraft movements.

Camera trap data and aircraft movement data were from the same sampling periods with aircraft movement data having been truncated to match camera trap deployment dates (between July 2019–May 2021). As hare strike data were available from 2012–2021 ($n = 238$), the entire data set was used to investigate the overlap patterns with aircraft movements and hare activity to ensure a sufficient sample size for robust estimates (Lashley et al., 2018; Meredith & Ridout, 2021). Truncating hare strike data to the sampling period dates (July 2019–May 2021) would result in an insufficient sample size ($n = 17$) to reliably estimate activity overlap.

Results

Camera trap detections

Of the potential 2,520 recording days (*i.e.*, 7 cameras recording for 360 calendar days), 2,144 were successful (85%). Camera failings were a result of removal by personnel, hares chewing through camera straps as well as battery and mechanical failure. A total of 684 independent mammalian detections were recorded on the airfield at Dublin Airport, from 5 species, across 360 calendar days (*i.e.*, full 24-hour periods; Table 1). The Irish hare was the most frequently detected species, making up 84.9% ($n = 574$) of detections, followed by the red fox (*Vulpes vulpes*) with 14.6% ($n = 100$) of detections. The European hedgehog (*Erinaceus europaeus*, $n = 5$), European rabbit (*Oryctolagus cuniculus*, $n = 3$) and domestic cat (*Felis catus*, $n = 2$) were rarely recorded. Over the course of the study period (360 days), a total of 114,559

aircraft movements were recorded at the airfield (Table 2).

Activity patterns

Hares demonstrated a largely crepuscular activity pattern, with peaks in activity recorded at, or close to, sunrise and sunset (Fig. 1). This pattern was less defined during the winter months, when hares were active over a longer period, likely due to the prolonged hours of darkness typical of the season in Ireland (*e.g.*, day length on winter solstice in Dublin is approximately 7 hours and 30 minutes, compared with approximately 17 hours on the summer solstice). In contrast, while aircraft movements were recorded continuously throughout the year, the majority of movements occurred according to a diurnal pattern (Fig. 1). Movements rapidly increased around 06:00 across all seasons and remained high for the duration of the day, decreasing in volume approaching midnight (00:00). Overlap between hares and aircraft was moderate across the year (58%; CI 53–60%) and across all seasons (Table 3), with spring having the highest activity overlap (63%; CI 55–65%). Activity overlap estimates for the sampling periods prior to the Covid-19 pandemic were slightly lower than those when both years were considered together (41–51%; Appendix 4).

Hare activity generally preceded aircraft activity, with a significant peak of activity at zero indicating contemporaneous activity patterns recorded only for the winter sampling period (peak = 0, $r = -0.576$, $t = 3.30$; Table 4), when hare activity was less confined to a strictly crepuscular pattern. Significant correlations between aircraft and hares were observed across all seasons, with activity patterns crossing zero throughout the year (peak = -3, $r = -0.628$, $t = 3.79$) as well as for the summer (peak = -1, $r = -0.592$, $t = 3.45$), autumn (peak = -2, $r = -0.608$, $t = 3.59$) and winter sampling periods (Table 4).

Table 1. Total number of seasonal detections of Irish hare and red fox using camera traps at Dublin Airport 2019–2021. Sampling commenced in the Summer of 2019 until Spring 2021.

Species	Year	Summer	Autumn	Winter	Spring	No. of independent triggers	No. days <i>in-situ</i>	No. of animal detections	No. of triggers per 50 days
Irish hare	1	40	30	45	59* [†]	174	152	175	57.2
<i>Lepus timidus hibernicus</i>	2	82*	46*	114*	158*	400	208	444	94.7
Red fox	1	18	9	10	12*	49	152	49	16.1
<i>Vulpes vulpes</i>	2	7*	13*	10*	21*	51	208	51	12.1

Year 1 = Summer 2019– Spring 2020.

Year 2 = Summer 2020– Spring 2021.

[†]Denotes the onset of the Covid-19 pandemic in Ireland.

*Denotes sampling periods during the Covid-19 pandemic in Ireland.

Table 2. Total number of Aircraft Movements (ACM) 2019–2021 recorded by the (daa) for the time frame during which the cameras were deployed on the airfield to record wildlife activity. Sampling commenced in the Summer of 2019 until Spring 2021.


Year	Summer	Autumn	Winter	Spring	Total	No. days	No. of ACM per 50 days	No. of ACM per 50 days prior to Covid-19
1	24,219	36,363*	17,287	2,363*†	80,232	152	26,392	32,445
2	16,035*	5,697*	8,022*	4,613*	34,367	211	8,144	NA

Year 1 = Summer 2019– Spring 2020.

Year 2 = Summer 2020– Spring 2021.

†Denotes the onset of the Covid-19 pandemic in Ireland.

*Denotes sampling periods during the Covid-19 pandemic in Ireland.

**Figure 1.** Overlap estimates of hare activity and aircraft movements at Dublin Airport. Shaded grey areas indicate times when activity overlapped. Dotted and solid black lines indicate sunrise/sunset times on the shortest day and longest day of the sample period respectively. Events are indicated along the x-axis for hares (blue) and aircraft movements (black).

Significant differences ($P < 0.001$) in hare activity relative to sunrise/sunset were recorded across all seasons at Dublin Airport ($F_{3,570} = 81.89$, $P < 0.0001$), with the exception of autumn–winter and spring–summer (Fig. 2). Due to disruptions caused by the Covid-19 pandemic to standard airfield activity patterns, interannual hare activity for each season was compared ($F_{3,566} = 37.46$, $P < 0.0001$) to ensure that activity patterns were a true

reflection of circadian activity with no significant differences observed between the same season for each year. Using Tukey's HSD test, small differences (diff) in mean activity offsets were observed for spring (diff: -0.87, 95% CI = -2.13, 0.39, $P = 0.42$), summer (diff: 1.32, 95% CI = -0.28, 2.92, $P = 0.19$), autumn (diff: -0.33, 95% CI = -2.27, 1.61, $P = 0.99$) and winter (diff: -0.62, 95% CI = -2.07, 0.84, $P = 0.90$).

Table 3. Annual and seasonal temporal overlap estimates for hare activity (2019–2021), aircraft movement activity (2019–2021) and reported strikes between hares and aircraft (2012–2021) at Dublin Airport with bootstrapped confidence intervals (95%).

Season	Overlap estimate (%)	Upper CI	Lower CI
i. Hare activity and aircraft activity temporal overlap (Fig. 1)			
Annual	58.3	53.3	59.9
Spring	62.7	54.8	64.7
Summer	55.0	45.5	59.7
Autumn	52.0	41.9	58.3
Winter	59.2	51.6	64.8
ii. Aircraft activity and recorded hare strike events (2012–2021) temporal overlap (Appendix 5)			
Annual	57.6	51.3	61.4
Spring	61.3	53.1	70.8
Summer	38.4	29.0	46.0
Autumn	62.6	52.9	73.0
Winter	61.7	52.3	68.0
iii. Hare activity and recorded hare strike events (2012–2021) temporal overlap (Fig. 3)			
Annual	85.8	79.5	90.3
Spring	66.1	54.2	74.0
Summer	74.5	64.2	84.5
Autumn	78.2	69.5	91.1
Winter	79.5	72.0	91.9

Table 4. Annual and seasonal associations and dissociations in temporal activity between hares and aircraft at Dublin Airport, estimated using Cross-Correlation Functions (CCF). r = Pearson's correlation coefficient.

Season	Hours			t -value	r
	Lag (from)	Lag (to)	Peak lag		
Annual	–12	–8	–9	3.14	0.557*
Annual	–5	0	–3	3.79	–0.628*
Spring	–5	–3	–4	2.42	–0.603*
Spring	–9	–9	–9	1.62	–0.327
Summer	–12	–8	–9	2.64	0.491*
Summer	–4	0	–1	3.45	–0.592*
Autumn	–11	–8	–9	2.28	0.438*
Autumn	–5	1	–2	3.59	–0.608*
Winter	–11	–8	–10	2.76	0.507*
Winter	–4	2	0	3.30	–0.576*
Winter	10	12	12	1.55	0.315

Negative lags indicate that hare activity preceded aircraft movements and positive lags indicate that hare activity followed aircraft movements. Zero indicates that activity was contemporaneous.

*Denotes significant lag.

Strike patterns

Strike events across the year followed a crepuscular pattern. The highest number of strikes was recorded simultaneously to peak hare activity in the morning (04:00–

07:59; 33% of strike events) and following the peak in the evening (21:00–23:59; 27% of strike events). Strike event times had a high ($\Delta > 0.75$) level of overlap with hare activity patterns across the year (86%; CI 80–90%, Fig. 3), as well as throughout the summer, autumn and winter seasons. Spring was the only season where a moderate overlap was recorded (66%; CI 54–74%).

Strike event times had a moderate overlap with aircraft activity times across the year (58%; CI 51–61%), with varying degrees of overlap across seasons. Highest overlap was recorded during the autumn period (62%; CI 53–73%) with the least amount of overlap recorded during the summer (39%; CI 29–46%; Table 3). Significant differences ($P < 0.001$) in strike events relative to sunrise/sunset were recorded ($F_{3,234} = 6.14$, $P < 0.0001$) for winter compared with all other seasons (Fig. 2). Aircraft movements generally picked up from approximately 06:00, with a high proportion of strike events recorded until 08:00 across the year. A second peak in strike events was recorded at night (21:00–23:59) when hourly aircraft movements were declining (Fig. 4, Appendix 5). Overall, 12% of strikes occurred between 07:00–07:59 and a further 12% between 23:00–23:59.

Discussion

Here we show that camera trap data can be used to identify the circadian and seasonal periods of increased strike risk in the air transportation sector (see Carswell et al., 2021). Prior research has investigated the suitability of camera traps for use on airfields for understanding species composition (Scheideman et al., 2017) and here we demonstrate their suitability in determining the activity patterns of a terrestrial mammal species in relation to aircraft movements. This study, using camera traps in a transportation management setting, represents a valuable addition to the growing literature on camera trap applications for wildlife conservation and management (e.g. Caravaggi et al., 2018; Garrote et al., 2019; Hofmeester et al., 2020; Jachowski et al., 2015; Schwartz et al., 2018). More importantly, it demonstrates the suitability of this survey method for quantifying ecological phenomena of management concern in busy, dynamic and heavily regulated environments.

Previous studies have demonstrated that lagomorphs alter activity patterns with increasing human influence on the landscape (Wong & Candolin, 2015; Ziege et al., 2016). However, the activity patterns of the Irish hare recorded during this study did not appear to be influenced by ongoing operations at Dublin Airport and were similar to previously recorded seasonal and circadian activity patterns for this sub-species (Caravaggi et al., 2018). The hare population at Dublin Airport




Figure 2. (A) Detection time of hare activity (2019–2021) on the airfield using motion-activated camera traps and (B) time of reported hare strike events (2012–2021) relative to sunrise and sunset for each season. Shaded areas indicate hours after sunset and lighter areas indicate hours after sunrise. Mean $\pm$ SD are represented by boxplots (left) and the density and spread of the data are represented by raincloud plots (right; Allen et al., 2021). The mean annual offset of events relative to sunrise and sunset across all seasons is represented by the dashed line.

exhibited a bimodal, crepuscular activity pattern, with clear peaks in activity at sunrise and sunset for the spring and summer seasons when days are longer. A higher rate of diurnal activity was also recorded during these seasons, likely due to energetic requirements and the need to continue foraging beyond the hours of (semi-) darkness. During the autumn and winter seasons, hares exhibited a trimodal activity pattern, with peaks at sunrise, sunset and at approximately midnight, as seen in some other mammal species (Brivio et al., 2016; Ikeda et al., 2019). Generally, activity times of the Irish hare at Dublin Airport preceded peak aircraft activity times, with winter being the only season when a significant peak in hare activity was contemporaneous with aircraft movements. Given that the majority of aircraft movements occurred in the daytime, this is to be expected as hares were predominantly active prior to the start of daily airfield operations.

The development of effective strike mitigation measures requires an understanding of strike events and factors driving these events. Overlap between hare activity and aircraft movements was moderate (58%) but was high between hare activity times and hare strike times at the airfield (85%), with strikes typically occurring according to a bimodal pattern with a peak at 04:00–07:59 and another at 21:00–23:59. Identifying times of higher risk is useful from a management execution and implementation perspective. These, for instance, are the times during which scaring and patrol efforts may be increased or aircraft may receive additional alerts about possible hare activity. Indeed, concentrating efforts for only 2 hours a day (07:00–07:59 and 23:00–23:59) could be greatly beneficial. However, as animals adjust activity seasonally with changing sunrise and sunset times, actual intraspecies activity peaks may change week to week based on natural and artificial light levels (Hoffmann et al., 2018). Indeed,




Figure 3. Overlap estimates of hare activity and hare strikes at Dublin Airport. Shaded grey areas indicate times when activity overlapped. Dotted and solid black lines indicate sunrise/sunset times on the shortest day and longest day of the sample period respectively. Events are indicated along the x-axis for hares (blue) and strikes (black).

we have demonstrated that hare activity at Dublin changes according to sunrise and sunset times and on a seasonal basis. Consequently, airfield managers need to operate a suite of mitigation measures that are flexible in their response to changing animal activity patterns.

Despite several road ecology studies demonstrating that increased volumes of road traffic are associated with increased roadkill rates (e.g. Haigh, 2012), strikes occurred at Dublin when aircraft movement numbers were relatively low, indicating that hare activity is a better indicator of strike risk than the volume of aircraft movements. This has important considerations should aircraft activity patterns be altered and suggests that strike risk may remain high at specific times of the day (periods of high hare activity) even during periods when aircraft activity patterns are altered (e.g. such as during the Covid-19 pandemic and economic crashes (Franke & John, 2011)). This furthermore demonstrates the

importance of maintaining high-quality strike data on a local and national scale (e.g. FAA, 2021).

With the onset of the Covid-19 pandemic, national lockdowns were implemented in Ireland at the end of March 2020 with aircraft movement numbers being severely impacted during the remainder of the year. As with road-traffic data on collisions with mammals (e.g. Łopucki et al., 2021), this reduction in air traffic coincided with a reduction in the number of hare strikes recorded at Dublin Airport. An average of 77% of recorded hare strikes at Dublin Airport occurred during April–December from 2012–2019. However, only 38% of strike events were recorded during this time frame in 2020. Although there was a small decrease (~8%) estimated in the size of the population once surveys resumed (June 2020–July 2021) from pre-lockdown data (SB, unpublished data), this is unlikely to explain the reduction in strike numbers. Despite this, annual strike rates (per




Figure 4. Activity times for the Irish hare (circadian), aircraft movements and recorded strike events (2012–2021) between hares and aircraft at Dublin Airport, across the year. Hare and aircraft activity data were collected between 2019–2021. Light grey shaded areas demonstrate the range of sunrise and sunset times through the study period (i.e., 21st June vs. 21st December). Dark grey shaded areas demonstrate hours of darkness across the whole study.

10,000 aircraft movements) with the Irish hare did not decrease, notwithstanding a 65% reduction in aircraft movements for 2020 compared with 2019 (daa, *unpublished data*). Indeed, strikes increased from 0.54 strikes per 10,000 aircraft movements in 2019 (Ball, Butler, et al., 2021a), to 0.92 strikes per 10,000 aircraft movements in 2020, indicating that the number of aircraft movements are at least partially responsible for the number of strike events. However, strike rates in 2019 were unusually low, potentially attributed to population management practices in 2018 (1.89 strikes per 10,000 aircraft movements; Ball, Butler, et al., 2021a) and habitat disturbance in 2019 due to ongoing construction works for a new runway. These observed changes in strike rate at the airfield could potentially be attributed to the consequential changes in aircraft movement patterns, land use changes by the hares or potentially due to increased naivety of the population (Mumme et al., 2000; Schwartz et al., 2020). Despite the conservation status of the Irish hare, strike events are unlikely to have population level impacts (Ball, Butler, et al., 2021a). Dublin Airport was not the only airfield to report an increased strike rate with hares, with Italian airports reporting an 81%

increase in hare strike rate during the 2020 lockdown periods (Montemaggiore, 2021).

Despite changes to air traffic volume at Dublin post the implementation of Covid-19 lockdown measures, the circadian activity of the hares at the airfield did not change between seasonal sampling periods (i.e., summer 2019 vs. summer 2020). While hare activity data for the spring prior to pandemic-related disruptions to aircraft traffic were not available, we do not believe that reduced aircraft movements impacted on hare activity at Dublin Airport. Circadian activity patterns for spring followed the same bimodal pattern as the summer sampling period and were consistent with previously published activities of the Irish hare (Caravaggi et al., 2018). While circadian activity of the hares was unchanged by the reduction of air traffic, camera traps documented a change in hare grouping behaviour at the airfield. Prior to the pandemic, more than one hare was recorded in a frame only 0.89% of the time and group size did not exceed two. Multiple hares were recorded 11% of the time during the same seasons in 2020, with up to four hares recorded within a single frame. Other mountain hare populations have been recorded to spend long periods of time under canopy

cover to avoid predation (Rehnus, 2014). However, by design, Dublin Airport is devoid of diverse habitat which could offer shelter to wildlife. With reports of increased and altered bird and wildlife activity at airfields with reduced air traffic (Schrimpf et al., 2021), the forming of these groups could be a social response to perceived increase in predation pressure at the airfield, with reduced aircraft movements to deter predators. Alternatively, reduced aircraft movements and consequential reduced disturbance could have allowed for the formation of leisurely social groups.

Daily and seasonal activities are thought to be two behavioural factors influencing mammal strike incidents with aircraft (Schwarz et al., 2014). As mammal strike events are becoming an increasing concern to the airline industry, understanding these patterns could help to mitigate strike risk at airfields by allowing for the targeted implementation of strike mitigation measures. Such measures could include increased runway patrols (e.g. Crain et al., 2015), species-specific noise and light harassment (Biondi et al., 2011), or giving prior warning to pilots operating during periods of increased risk. While aircraft movement temporal activity and volume are likely to play a role in strike patterns, we found wildlife activity itself to be more closely associated with strike patterns, highlighting the importance of understanding the ecology and life histories of the fauna using the airfield environment. Hence, while we use data for the Irish hare, these methods would be suitable for a cohort of terrestrial mammals associated with airfields (e.g. canids, Crain et al., 2015; ungulates, Biondi et al., 2011), assuming that an adequate number of detections (~100; Lashley et al., 2018) are obtained. High-quality data are vital to aid wildlife strike prevention research on airfields and for other modes of transportation within the sector (Steiner et al., 2014). An added benefit of data collection through the use of camera traps is the ability to capture the activities of sympatric species using one piece of equipment which can help inform risk and management decisions regarding multiple species (Appendix 6).

Conclusion

Using predator–prey data analytical methods and relatively inexpensive remote sensing equipment, we determined the activity patterns of a mammal species inhabiting the airfield in a large, international airport and identified periods of increased risk. Strike events were more closely associated with hare activity at the airfield rather than aircraft activity. This demonstrates the importance of identifying and understanding the wildlife populations utilising the airfield environment. These data can be used to inform the development of suitable mitigation strategies focussed on the species of concern (as opposed to the near-impossible task

of altering aircraft activity) and to identify periods of increased risk with other mammalian species at other airfields.

Acknowledgements

We gratefully acknowledge the contribution of the airfield staff and airport authorities at Dublin Airport for providing aircraft movement and hare strike data. Thanks to Gerry Keogh, Jim Eviston, Bob Navan and Clemence Parneix of the daa for project support and for providing bespoke data. Thank you to the fire crew at Dublin Airport for field assistance. Thank you to Thomas Kelly (UCC/daa) for the long-term collation of hare strike data and as a project mentor. To Allen Whitaker (UCC) for equipment support and technical guidance. This work was conducted as part of a PhD studentship funded by the Irish Research Council (IRC) and the daa in collaboration with University College Cork (project EBPPG/2018/43). Open access funding provided by IReL.

References

- Allen, M., Poggiali, D., Whitaker, K., Marshall, T.R., van Langen, J. & Kievit, R.A. (2021) Raincloud plots: a multi-platform tool for robust data visualization [version 2; peer review: 2 approved]. *Wellcome Open Research*, **4**, 63. <https://doi.org/10.12688/wellcomeopenres.15191.2>
- Altringer, L., Navin, J., Begier, M.J., Shwiff, S.A. & Anderson, A. (2021) Estimating wildlife strike costs at US airports: a machine learning approach. *Transportation Research Part D: Transport and Environment*, **97**, 102907. <https://doi.org/10.1016/j.trd.2021.102907>
- Arrondo, E., García-Alfonso, M., Blas, J., Cortes-Avizanda, A., De la Riva, M., Devault, T.L. et al. (2021) Use of avian GPS tracking to mitigate human fatalities from bird strikes caused by large soaring birds. *Journal of Applied Ecology*, **58**, 1411–1420. <https://doi.org/10.1111/1365-2664.13893>
- Askren, R.J., Dorak, B.E., Hagy, H.M., Eichholz, M.W., Washburn, B.E. & Ward, M.P. (2019) Tracking Canada geese near airports: using spatial data to better inform management. *Human-Wildlife Interact.*, **13**, 344–355. <https://doi.org/10.26077/yv1k-dg31>
- Ball, S., Butler, F., Caravaggi, A., Coughlan, N.E., Keogh, G., Callaghan, M.J.A.O. et al. (2021a) Hares in the long grass : increased aircraft related mortality of the Irish hare (*Lepus timidus hibernicus*) over a 30- year period at Ireland's largest civil airport. *European Journal of Wildlife Research*, **67–80**, 1–10. <https://doi.org/10.1007/s10344-021-01517-y>
- Ball, S., Caravaggi, A. & Butler, F. (2021b) Runway roadkill: a global review of mammal strikes with aircraft. *Mammal Review*, **1–16**, 420–435. <https://doi.org/10.1111/mam.12241>
- Biondi, K., Belant, J., Martin, J., DeVault, T. & Wang, G. (2011) White-tailed deer incidents with U.S. civil aircraft.

- Wildlife Society Bulletin*, **35**, 303–309. <https://doi.org/10.1002/wsb.46>
- Bolger, R. & Kelly, T. (2008) *Wildlife Management Plan 2008*. Dublin, Ireland: Dublin Airport Authority.
- Brivio, F., Bertolucci, C., Tettamanti, F., Filli, F., Apollonio, M. & Grignolio, S. (2016) The weather dictates the rhythms: alpine chamois activity is well adapted to ecological conditions. *Behavioral Ecology and Sociobiology*, **70**, 1291–1304. <https://doi.org/10.1007/s00265-016-2137-8>
- Caravaggi, A., Gatta, M., Vallely, M., Hogg, K., Freeman, M., Fadaei, E. et al. (2018) Seasonal and predator-prey effects on circadian activity of free-ranging mammals revealed by camera traps. *Peer J*, **6**, 1–27. <https://doi.org/10.7717/peerj.5827>
- Caravaggi, A., Montgomery, I. & Reid, N. (2017) Management and control of invasive brown hares (*Lepus europaeus*): contrasting attitudes of selected environmental stakeholders and the wider rural community. *Biology Environment*, **117B**, 53–63. <https://doi.org/10.3318/BIOE.2017.08>
- Caravaggi, A., Zaccaroni, M., Riga, F., Schai-Braun, S.C., Dick, J.T.A., Montgomery, W.I. et al. (2016) An invasive-native mammalian species replacement process captured by camera trap survey random encounter models. *Remote Sensing in Ecology and Conservation*, **2**, 45–58. <https://doi.org/10.1002/rse2.11>
- Carswell, B.M., Rea, R.V., Searing, G.F. & Hesse, G. (2021) Towards building a species-specific risk model for mammal-aircraft strikes. *Journal of Air Transport Management*, **15**, 288–303.
- Clair, C.C.S., Backs, J., Friesen, A., Gangadharan, A., Gilhooly, P., Murray, M. et al. (2019) Animal learning may contribute to both problems and solutions for wildlife-train collisions. *Philosophical Transactions of the Royal Society B*, **374**, 20180050. <https://doi.org/10.1098/rstb.2018.0050>
- Crain, A., Belant, J. & DeVault, T. (2015) Carnivore incidents with U.S. civil aircraft. *Transportation Research Part D*, **36**, 160–166. <https://doi.org/10.1016/j.trd.2014.12.001>
- Dolbeer, R. & Begier, M.J. (2021) *Strikes to civil aircraft in the United States, 1990–2019*. Washington, DC: Federal Aviation Administration. U.S Department of Agriculture, Animal and Plant Health Inspection Service, Wildlife Services.
- Dorsey, B.P., Clevenger, A. & Rew, L.J. (2017) Relative risk and variables associated with bear and ungulate mortalities along a railroad in the Canadian Rocky Mountains. In: Borda-de-Água, L., Barrientos, R., Beja, P. & Pereira, H.M. (Eds.) *Railway Ecology*. Cham, Switzerland: Springer, pp. 135–155.
- FAA. (2021) *National wildlife strike database*. Washington, DC: Federal Aviation Administration. <https://wildlife.faa.gov/home>
- Franke, M. & John, F. (2011) What comes next after recession? - Airline industry scenarios and potential end games. *Journal of Air Transport Management*, **17**, 19–26. <https://doi.org/10.1016/j.jairtraman.2010.10.005>
- Garrote, G., Pérez De Ayala, R., Álvarez, A., Martín, J.M., Ruiz, M., De Lillo, S. et al. (2019) Improving the random encounter model method to estimate carnivore densities using data generated by conventional camera-trap design. *Oryx*, **1–6**, 99–104. <https://doi.org/10.1017/S0030605318001618>
- Haigh, A. (2012) Annual patterns of mammalian mortality on Irish roads. *Hystrix*, **23**, 58–66. <https://doi.org/10.4404/hystrix-23.2-4747>
- Hoffmann, J., Palme, R. & Eccard, J.A. (2018) Long-term dim light during nighttime changes activity patterns and space use in experimental small mammal populations. *Environmental Pollution*, **238**, 844–851. <https://doi.org/10.1016/j.envpol.2018.03.107>
- Hofmeester, T.R., Young, S., Juthberg, S., Singh, N.J., Widemo, F., Andrén, H. et al. (2020) Using by-catch data from wildlife surveys to quantify climatic parameters and timing of phenology for plants and animals using camera traps. *Remote Sensing in Ecology and Conservation*, **6**, 129–140. <https://doi.org/10.1002/rse2.136>
- Ikedo, T., Takahashi, H., Igota, H., Matsuura, Y., Azumaya, M., Yoshida, T. et al. (2019) Effects of culling intensity on diel and seasonal activity patterns of sika deer (*Cervus nippon*). *Scientific Reports*, **9**, 1–8. <https://doi.org/10.1038/s41598-019-53727-9>
- Jachowski, D.S., Katzner, T., Rodrigue, J.L. & Ford, W.M. (2015) Monitoring landscape-level distribution and migration Phenology of Raptors using a volunteer camera-trap network. *Wildlife Society Bulletin*, **39**, 553–563. <https://doi.org/10.1002/wsb.571>
- Jaren, V., Andersen, R., Ulleberg, M., Pedersen, P. & Wiseth, B. (1991) Moose-train collisions: the effects of vegetation removal with a cost-benefit analysis. *Alces*, **27**, 93–99.
- Kelly, T.C., Sleeman, P., Coughlan, N.E., Dillane, E. & Callaghan, M.J. (2017) Bat collisions with civil aircraft in the Republic of Ireland over a decade suggest negligible impact on aviation safety. *European Journal of Wildlife Research*, **63**, 23–26. <https://doi.org/10.1007/s10344-017-1081-x>
- Kitowski, I. (2016) The impact on aviation operations at Polish civil airfields caused by mammals. *Bezpieczeństwo i Technika Pożarnicza*, **42**, 57–63. <https://doi.org/10.12845/bitp.42.2.2016.5>
- Laist, D.W., Knowlton, A.R., Mead, J.G., Collet, A.S. & Podesta, M. (2001) Collisions between ships and whales. *Marine Mammal Science*, **17**, 35–75. <https://doi.org/10.1111/j.1748-7692.2001.tb00980.x>
- Lashley, M.A., Cove, M.V., Chitwood, M.C., Penido, G., Gardner, B., Deperno, C.S. et al. (2018) Estimating wildlife activity curves: comparison of methods and sample size. *Scientific Reports*, **8**, 1–11. <https://doi.org/10.1038/s41598-018-22638-6>
- Lendrum, P.E., Crooks, K.R. & Wittemyer, G. (2017) Changes in circadian activity patterns of a wildlife community post

- high-intensity energy development. *Journal of Mammalogy*, **98**, 1265–1271. <https://doi.org/10.1093/jmammal/gyx097>
- Lopucki, R., Kitowski, I., Perlińska-Teresiak, M. & Klich, D. (2021) How is wildlife affected by the covid-19 pandemic? Lockdown effect on the road mortality of hedgehogs. *Animals*, **11**, 1–8. <https://doi.org/10.3390/ani11030868>
- Meijer, J.H., Michel, S., VanderLeest, H.T. & Rohling, J.H.T. (2010) Daily and seasonal adaptation of the circadian clock requires plasticity of the SCN neuronal network. *The European Journal of Neuroscience*, **32**, 2143–2151. <https://doi.org/10.1111/j.1460-9568.2010.07522.x>
- Meredith, M. & Ridout, M. (2021) *Overlap: estimates of coefficient of overlapping for animal activity patterns*. <https://cran.r-project.org/web/packages/overlap/overlap.pdf>
- MET Eireann. (2021). Dublin airport monthly data [WWW Document]. <https://www.met.ie/climate/available-data/monthly-data> (accessed 9.20.21).
- Montemaggiore, A. (2021) Wildlife strike hazard and lockdown at Italian airports. In: *Proceedings of the Presented at the World Bird Strike Virtual Conference*. Virtual: World Birdstrike Association.
- Monterroso, P., Alves, P.C. & Ferreras, P. (2014) Plasticity in circadian activity patterns of mesocarnivores in Southwestern Europe: Implications for species coexistence. *Behavioral Ecology and Sociobiology*, **68**, 1403–1417. <https://doi.org/10.1007/s00265-014-1748-1>
- Mumme, R.L., Schoech, S.J., Woolfenden, G.E. & Fitzpatrick, J.W. (2000) Life and death in the fast lane: demographic consequences of road mortality in the Florida Scrub-Jay. *Conservation Biology*, **14**, 501–512. <https://doi.org/10.1046/j.1523-1739.2000.98370.x>
- Pereira, J.A. (2010) Activity pattern of Geoffroy's cats (*Leopardus geoffroyi*) during a period of food shortage. *Journal of Arid Environments*, **74**, 1106–1109. <https://doi.org/10.1016/j.jaridenv.2010.03.017>
- Phillips, A.J.K., Fulcher, B.D., Robinson, P.A. & Klerman, E.B. (2013) Mammalian rest/activity patterns explained by physiologically based modeling. *PLoS Computational Biology*, **9**, e1003213. <https://doi.org/10.1371/journal.pcbi.1003213>
- Pokorny, B., Cerri, J. & Bužan, E. (2022) Wildlife roadkill and COVID-19: a biologically significant, but heterogeneous, reduction. *Journal of Applied Ecology*, **59**(5) 1291–1301.
- Popp, J.N. & Boyle, S.P. (2017) Railway ecology: Underrepresented in science? *Basic and Applied Ecology*, **19**, 84–93. <https://doi.org/10.1016/j.baae.2016.11.006>
- R Core Team. (2021) *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Rehnus, M. (2014) The 24-hour cycle of the mountain hare *Lepus timidus* Linnaeus, 1758. *Zoological Garten*, **83**, 140–145. <https://doi.org/10.1016/j.zoolgart.2014.10.001>
- Reid, N., McDonald, R.A. & Montgomery, W.I. (2010) Homogeneous habitat can meet the discrete and varied resource requirements of hares but may set an ecological trap. *Biological Conservation*, **143**, 1701–1706. <https://doi.org/10.1016/j.biocon.2010.03.041>
- Ross, J., Hearn, A.J., Johnson, P.J. & Macdonald, D.W. (2013) Activity patterns and temporal avoidance by prey in response to Sunda clouded leopard predation risk. *Journal of Zoology*, **290**, 96–106. <https://doi.org/10.1111/jzo.12018>
- Scheideman, M., Rea, R., Hesse, G., Soong, L., Green, C., Sample, C. et al. (2017) Use of wildlife camera traps to aid in wildlife management planning at airports. *Journal of Air Transport Management*, **11**, 408–419.
- Schrimpf, M.B., des Brisay, P.G., Johnston, A., Smith, A.C., Sánchez-Jasso, J., Robinson, B.G. et al. (2021) Reduced human activity during COVID-19 alters avian land use across North America. *Science Advances*, **7**, 1–12. <https://doi.org/10.1126/sciadv.abf5073>
- Schwartz, A.L., Shilling, F.M. & Perkins, S.E. (2020) The value of monitoring wildlife roadkill. *European Journal of Wildlife Research*, **66**(1), 1–12. <https://doi.org/10.1007/s10344-019-1357-4>
- Schwartz, A.L., Williams, H.F., Chadwick, E., Thomas, R.J. & Perkins, S.E. (2018) Roadkill scavenging behaviour in an urban environment. *Journal of Urban Ecology*, **4**(1), juy006. <https://doi.org/10.1093/jue/juy006>
- Schwarz, K., Belant, J., Martin, J., DeVault, T. & Wang, G. (2014) Behavioral traits and airport type affect mammal incidents with U.S. civil aircraft. *Environmental Management*, **54**, 908–918. <https://doi.org/10.1007/s00267-014-0345-4>
- Steiner, W., Leisch, F. & Hackländer, K. (2014) A review on the temporal pattern of deer-vehicle accidents: impact of seasonal, diurnal and lunar effects in cervids. *Accident; Analysis and Prevention*, **66**, 168–181. <https://doi.org/10.1016/j.aap.2014.01.020>
- UKCAA Safety Regulation Group CAP. (2008) *CAP 772 Birdstrike Risk Management for Aerodromes*. London, UK.
- Viviano, A., Mori, E., Fattorini, N., Mazza, G., Lazzeri, L., Panichi, A. et al. (2021) Spatiotemporal overlap between the European brown hare and its potential predators and competitors. *Animals*, **11**, 562. <https://doi.org/10.3390/ani11020562>
- Wolfe, A. & Hayden, T.J. (1996) Home range sizes of Irish mountain hares on coastal grassland. *Biological Environment*, **96**, 141–146.
- Wong, B.B.M. & Candolin, U. (2015) Behavioral responses to changing environments. *Behavioral Ecology*, **26**, 665–673. <https://doi.org/10.1093/beheco/aru183>
- Wright, P.G.R., Coomber, F.G., Bellamy, C.C., Perkins, S.E. & Mathews, F. (2020) Predicting hedgehog mortality risks on British roads using habitat suitability modelling. *PeerJ*, **2020**, 1–22. <https://doi.org/10.7717/peerj.8154>
- York, D.L., Cummings, J.L. & Wedemeyer, K.L. (2000) Movements and distribution of radio-collared Canada geese in Anchorage, Alaska. *Northwest Nature*, **81**, 11. <https://doi.org/10.2307/3536894>

- Zanni, M., Brivio, F., Grignolio, S. & Apollonio, M. (2021) Estimation of spatial and temporal overlap in three ungulate species in a Mediterranean environment. *Mammal Research*, **66**, 149–162. <https://doi.org/10.1007/s13364-020-00548-1>
- Ziege, M., Babitsch, D., Brix, M., Kriesten, S., Straskraba, S., Wenninger, S. et al. (2016) Extended diurnal activity patterns of European rabbits along a rural-to-urban gradient. *Mammalian Biology*, **81**, 534–541. <https://doi.org/10.1016/j.mambio.2016.07.002>

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Appendix 1. Images demonstrating that camera traps were effective at capturing activity during hours of daylight (*top*) and darkness (*bottom*).

Appendix 2. Minimum Metadata Standards for camera trap deployment. This is in in Excel format attached separately.

Appendix 3. camera trap deployment log.

Appendix 4. Overlap estimates of hare and aircraft activity for sampling periods prior to the Covid-19 pandemic.

Appendix 5. Overlap estimates of aircraft activity and recorded hare strike events (2012–2021) temporal overlap.


Appendix 6. An example of how the methodology is applicable to other terrestrial mammal species.

Hareport hazard: Identifying hare activity patterns and increased mammal-aircraft strike risk at an International Airport

Samantha Ball, Anthony Caravaggi, Fidelma Butler

Supplementary material

Appendix 1-Images demonstrating that camera traps were effective at capturing activity during hours of daylight (*top*) and darkness (*bottom*)



Appendix 2- Minimum Metadata Standards for camera trap deployment. This is in in Excel format attached separately.

Appendix 3- camera trap deployment log

Table S.1. Camera trap deployment dates for each sampling period


Year	Season	Deployment date from	Deployment date to	No. of days
2019	Summer	09/07/2019	10/08/2019	32
2019	Autumn	27/09/2019	22/11/2019	56
2020	Winter	10/01/2020	11/02/2020	32
2020	Spring*	29/04/2020	31/05/2020	32
2020	Summer*	17/06/2020	25/08/2020	69
2020	Autumn*	20/10/2020	30/11/2020	41
2020	Winter*	03/12/2020	28/01/2021	56
2021	Spring*	08/04/2021	20/05/2021	42
Total:				360

Appendix 4- Overlap estimates of hare and aircraft activity for sampling periods prior to the Covid-19 pandemic


Table S.2. Seasonal temporal overlap estimates for hare activity and aircraft movement activity for the three recording seasons prior to the Covid-19 pandemic with bootstrapped confidence intervals (95%).

Season	Overlap estimate (%)	Upper CI (%)	Lower CI (%)
Summer 2019	47.8	33.5	56.1
Autumn 2019	41.4	26.0	50.6
Winter 2019-2010	50.8	35.6	59.5


Summer 2019




Autumn 2019




Winter 2020



Appendix 5- Overlap estimates of aircraft activity and recorded hare strike events (2012-2021) temporal overlap



Appendix 6- An example of how the methodology is applicable to other terrestrial mammal species



S4. Overlap estimates of fox (*Vulpes vulpes*) activity and aircraft activity at Dublin Airport. Shaded grey areas indicate times when activity overlapped.

Column name	Form Value	Definition	Data Type
project_name	Project Name	A short name that uniquely identifies the project dataset and relates to project code (400 chars)	String
project_id	Project ID	identification for the project. It should be a relatively short	String
project_short_name	Project Short Name	A short name for you project that can only be 43	String
project_objectives	Project Objectives	What is your study objective?	String
project_species	Project Species	Is your study objective to monitor or	String
project_species_individual	Project Species Individual	If individual, which species? List the species	String
project_sensor_layout	Project Sensor Layout	What is the sensor layout or sampling	String
project_sensor_layout_targeted_type	Project Sensor Layout Targeted Type	what wildlife sign or other	String
project_bait_use	Project Bait Use	Were sensors baited at all or some of my	String
project_bait_type	Project Bait Type	If yes, what bait?	String
project_stratification	Project Stratification	Are your sensors	String
project_stratification_type	Project Stratification Type	If yes, list the strata	String
project_sensor_method	Project Sensor Method	sampling based on motion sensor	String
project_individual_animals	Project Individual Animals	Will marked individuals (or	String
project_blank_images	Project Blank Images	Were blank images removed from this dataset?	String
project_sensor_cluster	Project Sensor Cluster	sensors setup in pairs or	String
project_admin	Project Admin	The name of the project administrator	String
project_admin_email	Project Admin Email	Project administrators email-- 255	String
country_code	Country Code	3-letter ISO code	String

embargo	Embargo Period	Length of embargo in months	Integer
metadata_license	License Metadata	The license you would like associatd for the project	String
image_license	License Images	The license you would like associated to all of the media (currently images but will include other media types in the future) that you will be uploading into Wildlife Insights.	String
project_type	Project Type	Defines whether identification occured at the image or sequence level.	enum

Values List	Default Value	Value
None	None	Circadian activity at Dublin Airport
None	None	Circadian activity at Dublin Airport
None	None	Circadian activity at Dublin Airport
None	None	To identify the circadian activity of a resident population of Irish hare at the airfield at Dublin Airport
Individual, Multiple	Multiple	Irish hare (<i>Lepus timidus hibernicus</i>)
Multi-select	None	
Systematic, Randomized, Convenience,	Systematic	
	None	
Yes, Some, No	No	
Meat, Visual, Acoustic, Other	None	
Yes, No	No	
List the strata (free text)	None	
Detection, Time lapse, Both	Sensor Detection	Motion sensor
Yes, No	No	
Yes, Some, No	No	Yes
Yes, No	No	No
None	None	Samantha Ball
None	None	samantha.ball@ucc.ie
Three letter ISO codes are	None	IRL

0-48	0	
CC0, CC-BY	CC0	
CC0, CC-BY, CC-BY-NC	CC0	
Image,Sequence	Image	Image

Column name	Form Value	Definition	Data Type	Values List
project_id	Project ID	An internal identifier	String	None
camera_id	Camera IDs	An internal identifier	String	None
make	Make	Manufacturer of the camera	String	None
model	Model	The model of the camera	String	None
serial_number	Serial Number	The serial number of the camera	String	None
year_purchased	Year Purchased	The date the camera was purchased	Date (yyyy-mm-dd)	None

Default Value	Value
None	Circadian activity at Dublin Airport
None	A, B,C,D,E,F,G
None	Bushnell
None	Trophy Cam HD (model 119476)
None	12348571
None	Unknown

Column name	Form Value
project_id	Project ID
deployment_id	Deployment ID
subproject_name	Subproject name
subproject_design	Subproject design
placename	Deployment Location ID
longitude	Longitude
latitude	Latitude
start_date	Camera Deployment Begin Date
end_date	Camera Deployment End Date
event_name	Event
event_description	Event Description
event_type	Event Type
bait_type	Bait Type
bait_description	Bait Description
feature_type	Feature Type
feature_type_methodology	Feature Type Methodology
camera_id	Camera ID
quiet_period	Quiet Period Setting
camera_functioning	Camera Failure Details
sensor_height	Height
height_other	Height Other
sensor_orientation	Angle
orientation_other	Angle Other
recorded_by	Recorded By
plot_treatment	Plot Treatment
plot_treatment_description	Plot treatment description
detection_distance	Detection Distance

Definition	Default Value
should be a relatively short name that makes the	None
(within the project)	None
within a project. Subprojects are similar to Events	None
Text field to describe subproject design methods	None
deployment (within the project)	None
degrees using datum WGS84. Minimum 4	None
degrees using datum WGS84. Minimum 4	None
recorded observations. This also marks the date a	None
the SD card was retrieved. If the exact timestamp	None
'events' (not to be confused with events of	None
A description that defines the events.	None
example, some projects may put sensors out	None
standardized list with No Bait and Other Bait	None
Bait type if necessary. This field should be left	None
is associated with	None
	None
Camera ID (see Cameras tab)	None
activity in the sensor will not trigger the shutter.	0
deployment; standardized list	Camera Functioning
sensor? If user selection = "other", please	Knee height
If other, please describe	
What angle were generally used to deploy the sensor?	Parallel
If other, please describe	None
the person installing the camera	None
A parcel of land defined by a specific function, property, or purpose. Examples include types of agriculture and different stages of controlled burns.	None or NULL
General description of the plot treatment	None or NULL
Maximum distance at which a camera triggered, as tested during deployment, measured in meters	NULL

Value
Circadian activity at Dublin Airport
Circadian activity at Dublin Airport
Circadian activity at Dublin Airport
2021)
Dublin Airport, South Perimeter Road
-6.269137
53.421585
See tab 'deployment dates'
See tab 'deployment dates'
spring, summer, autumn, winter
Cameras were set up and allocated a sampling year (1 or 2), a season, and location (A-> G)
Season and year
None
Permanent structure present on the airfield (e.g. gate post) to allow for safe camera attachment
A,B,C,D,E,F,G
30 seconds between triggers
Knee height
~15 degrees downwards tilt
Samantha Ball
Airport grasslands



Year	Season	Deployment date from	Deployment date to	No. of days
2019	Summer	7/9/2019	8/10/2019	32
2019	Autumn	9/27/2019	11/22/2019	56
2020	Winter	1/10/2020	2/11/2020	32
2020	Spring*	4/29/2020	5/31/2020	32
2020	Summer*	6/17/2020	8/25/2020	69
2020	Autumn*	10/20/2020	11/30/2020	41
2020	Winter*	12/3/2020	1/28/2021	56
2021	Spring*	4/8/2021	5/20/2021	42
Total:				360

Column name	Form Value	Definition
project_id	Project ID	See Project
deployment_id	Deployment ID	See Deployment
image_id	Image ID	Unique identifier
location	Location	Location of the
identified_by	Photo Type Identified by	First and last name
wi_taxon_id	wi_taxon_identifier	A permanent ID
class	Class	Latin name for
order	Order	Latin name for
family	Family	Latin name for
genus	Genus	Latin name for
species	Species	Latin name for
common_name	Species Common Name	Common name
uncertainty	Uncertainty	The uncertainty
timestamp	Date_Time Captured	Date and time
number_of_objects	Number of Objects	Number of objects
highlighted	Image Favorite	Mark as a good
age	Age	Age of animal
sex	Sex	Sex of animal
animal_recognizable	Animal recognizable	Was the animal
individual_id	Individual ID	Individual animal
individual_animal_notes	Individual Image and Animal Notes	Notes about the
markings	Markings	Distinct markings
external_sequence_id	External Sequence ID	Unique identifier
sequence_start_time	External Sequence Start Time	The start date t



Data Type	Values List	Default Value
String	None	None
String	None	None
String or number	None	None
String	None	None
String	None	None
String	Reference WI Global Taxonomy table.	A valid wi_taxon_identifier
String	Reference WI Global Taxonomy table.	If present a valid Class from the WI Global Taxonomy table.
String	Reference WI Global Taxonomy table.	If present a valid Class from the WI Global Taxonomy table.
String	Reference WI Global Taxonomy table.	If present a valid Class from the WI Global Taxonomy table.
String	Reference WI Global Taxonomy table.	If present a valid Class from the WI Global Taxonomy table.
String	Reference WI Global Taxonomy table.	If present a valid Class from the WI Global Taxonomy table.
String	Reference WI Global Taxonomy table.	If present a valid common name from the WI Global Taxonomy table.
String	Absolutely sure	Don't know
Date (yyyy-mm-dd)	None	None
Numeric	Between 1-175	1
Integer	1,0	0
String	Adult, Juvenile,	Unknown
String	Male, Female,	Unknown
String	Yes, No	No
String	None	None
String	None	None
String	None	None
String	None	None or NULL
Date (yyyy-mm-dd)	None	NULL

[illegible]

Density of Activity



Density of Activity



Density of Activity

